

generic for phenergan medication

[\[PDF\] buy cialis in johor bahru](#)

[\[PDF\] viagra italy buy](#)

[\[PDF\] topamax generic effectiveness](#)

[\[PDF\] viagra verpackung original](#)

[\[PDF\] tramadol sales](#)

[\[PDF\] where to buy viagra in pattaya](#)

[\[PDF\] how much does 20 percocet cost without insurance](#)

Follow all directions on your prescription label. It is not known whether Phenergan will harm an unborn baby. Sedation trazodone , promethazine , lorazepam , hydroxyzine , Ativan , fentanyl , More Phenergan is used to treat allergy symptoms such as itching, runny nose, sneezing, itchy or watery eyes, hives, and itchy skin rashes. Light Sedation promethazine , diazepam , Valium , midazolam , chlordiazepoxide , chlorpromazine , More Stop using Phenergan and call your doctor at once if you have twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs. Get up slowly and steady yourself to prevent a fall. Call your doctor for medical advice about side effects. Skip the missed dose if it is almost time for your next scheduled dose. Your doctor may occasionally change your dose to make sure you get the best results. Phenergan can cause side effects that may impair your thinking or reactions. Phenergan is not for use in treating symptoms of asthma, pneumonia, or other lower respiratory tract infections. Do not start using a new medication without telling your doctor. If a child is using this medicine, tell your doctor if the child has any changes in weight. We comply with the HONcode standard for trustworthy health information - verify here. Greater than or equal to 2 years: Be careful if you drive or do anything that requires you to be alert. Learn about Phenergan (Promethazine) may treat, uses, dosage, side effects, drug interactions, warnings, patient labeling, reviews, and related medications. Phenergan (promethazine) is used to treat allergy symptoms and prevent motion sickness. Includes Generic Name: promethazine (oral) (pro METH a zeen) What is Phenergan? Phenergan (promethazine) belongs to a group of drugs called phenothiazines. It works by changing the actions of chemicals in your brain. ?Dosage ?Phenergan (FDA) ?Phenergan (Promethazine ?Side Effects. Phenergan official prescribing information for healthcare professionals. Generic Name: promethazine hydrochloride . The management of NMS should include 1) immediate discontinuation of promethazine HCl, antipsychotic drugs, if any, and other drugs not essential to concurrent therapy, 2) intensive symptomatic. Consumer information about the medication PROMETHAZINE - ORAL (Phenergan), includes side effects, drug interactions, recommended dosages, and storage information. Read more about the prescription drug PROMETHAZINE - ORAL. GENERIC NAME: PROMETHAZINE - ORAL (proe-METH-a-zeen). Drug information on Pentazine, Phenergan (promethazine (oral)), includes drug pictures, side effects, drug interactions, directions for use, symptoms of overdose, Generic Name: promethazine (oral) (Pronunciation: pro METH a zeen) Promethazine may also be used for other purposes not listed in this medication guide. Quick facts about Phenergan. Drug class: Gastrointestinal; Rx status: Prescription only; Generic status: Lower-cost generic available (promethazine). Upsides. Available in generic. Available as a syrup or tablet that you can take by mouth, or as a rectal suppository if you're unable to take anything by mouth. People's. Phenergan Oral tablet drug summary. Find medication information including related drug classes, side effects, patient statistics and answers to frequently asked questions. Visit unahistoriafantastica.com for more details. Jump to Generic - Promethazine. For more information on this medication choose from the list of selections below. Overview Black Box Warning Uses Brand Names Drug Class Side Effects Drug Interactions Drug Precautions Food Interactions Inform MD Pregnancy Lactation Drug Usage Drug Dosage. Promethazine is a neuroleptic medication and first-generation antihistamine of the phenothiazine family. The drug has strong sedative and weak antipsychotic effects. It also reduces motion sickness and has antiemetic (Histamine H1 receptors but it does not block the production of histamine) and anticholinergic properties. Promethazine hydrochloride should not be used in pediatric patients less than 2 years old because of the potential for fatal respiratory depression. Exercise caution when administering to pediatric patients 2 years or older and use at the lowest effective dose. Avoid concomitant use of other drugs with respiratory depressant.