

pharmacologic category of atorvastatin

[\[PDF\] fair price for vicodin](#)

[\[PDF\] purchase viagra online from canada](#)

[\[PDF\] etoricoxib pharmacokinetics](#)

[\[PDF\] is advil the same as generic ibuprofen](#)

[\[PDF\] buy nolvadex research](#)

[\[PDF\] cephalixin suspension price](#)

[\[PDF\] where to buy generic cialis safely](#)

Also contraindicated in pregnant or breast-feeding women and in women of childbearing age except those with no risk of becoming pregnant. Side effects in more detail. Mean plasma elimination half-life of atorvastatin is about 14 hours, but the half-life of inhibitory activity for HMG-CoA reductase is 20 to 30 hours because of the contribution of active metabolites. Treatment of primary dysbetalipoproteinemia Type III not adequately responsive to diet. Adults and children older than age 9: To reduce risk of MI and stroke in patients with type 2 diabetes and at risk see full labeling. History of liver disease. Tell patient to take drug with liquid other than grapefruit juice. Elevates drug levels and increases risk of adverse effects. This is not a complete list of side effects and others may occur. Take the medicine at the same time each day. Take Lipitor exactly as prescribed by your doctor. Discontinue if myopathy or elevated CPK levels occur. LIPITOR (Atorvastatin) drug information & product resources from MPR including dosage information, educational materials, & patient assistance. Adjunct to diet in primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia (Types IIa and IIb) to reduce total-C, LDL-C, apo B and TG, and atorvastatin calcium. Lipitor. Pharmacologic classification: 3-hydroxymethylglutaryl-coenzyme A (HMG-CoA) reductase inhibitor. Therapeutic classification: antilipemic. Pregnancy risk category X. Brand Names: U.S.. Lipitor. Pharmacologic Category. Antilipemic Agent, HMG-CoA Reductase Inhibitor. Pharmacology. Inhibitor of 3-hydroxymethylglutaryl coenzyme A (HMG-CoA) reductase, the rate-limiting enzyme in cholesterol synthesis (reduces the production of mevalonic acid from HMG-CoA); this then results in a. Atorvastatin, marketed under the trade name Lipitor among others, is a member of the medication class known as statins, which are used primarily as a lipid-lowering agent and for prevention of events associated with cardiovascular disease. Like all statins, atorvastatin works by inhibiting HMG-CoA reductase, an enzyme. Atorvastatin initiated before the ninth week post-transplant (n = 33, mean age +/- years) was compared to a control group who did not receive the drug inhibitors; rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria have been reported with atorvastatin and other drugs in this class. Dyslipidemia (Fredrickson Types IIa and IIb). LIPITOR reduces total-C, LDL-C, VLDL-C, apo B, and TG, and increases HDL-C in patients with hypercholesterolemia and mixed dyslipidemia. Therapeutic response is seen within 2 weeks, and maximum response is usually achieved within 4 weeks and maintained during. Jul 31, - Learn about Lipitor (Atorvastatin Calcium) may treat, uses, dosage, side effects, drug interactions, warnings, patient labeling, reviews, and related In adult patients with type 2 diabetes, and without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as. General. Pronunciation: a-tore-va-stat-in/e-zet-i-mibe. Trade Name(s). Lipitruzet. Pregnancy Category Category X (for atorvastatin) Category C (ezetimibe). Ther. Class. lipid-lowering agents. Pharm. Class. hmg coa reductase inhibitors cholesterol absorption inhibitors. Primary hypercholesterolemia and mixed dyslipidemia. Indicated as an adjunct to diet for treatment of elevated total-C, Apo B, and TG levels and to increase HDL-C in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia (Fredrickson type IIa and IIb); mg PO qDay. Drug information on Lipitor (atorvastatin), includes drug pictures, side effects, drug interactions, directions for use, symptoms of overdose, and what to avoid. Atorvastatin is used to treat high cholesterol, and to lower the risk of stroke, heart attack, or other heart complications in people with type 2 diabetes, coronary heart.