

is paxil used to treat bipolar disorder

[\[PDF\] makers of generic wellbutrin](#)

[\[PDF\] what pharmaceutical company makes femara](#)

[\[PDF\] cost lipitor vs atorvastatin](#)

[\[PDF\] nitroglycerin in labor and delivery](#)

[\[PDF\] clindamycin and benzoyl peroxide cost](#)

[\[PDF\] mirtazapine prescription](#)

[\[PDF\] how much is generic strattera](#)

These medicines treat full episodes of mania or depression that last for several days or weeks at a time. Which Bipolar Medicine Is Best? If you have sleep problems along with bipolar symptoms, you may get a type of drug called a benzodiazepine. They also help to keep your moods from interfering with work, school, or your social life. Take it at the same time every day. Many side effects will go away after a few weeks of treatment. Stick to Your Treatment Medications for bipolar disorder are powerful drugs, and you must take them exactly as your doctor recommends. Medicines for Bipolar Depression Most of the time, doctors will start bipolar treatment by prescribing a mood-stabilizing drug like lithium. Keep in mind that the term "mood stabilizer" can be misleading. Because of this risk, your doctor should keep track of you closely if you take one. Nausea Tremors Hair loss Sexual problems Weight gain Liver damage Kidney damage Diarrhea Belly pain Skin reaction Some medications can affect how well your liver works or the amount of white blood cells or platelets you have. You may need to try several different kinds and different doses to figure out the right approach. Similarly, wait two weeks after stopping Paxil before starting to take any MAO inhibitor. After two weeks have passed, start Paxil at a low dosage and increase gradually until an effective level is reached. Continued Side Effects of Bipolar Drugs Like any drug, bipolar medicines can cause some side effects. A weekly pillbox can help you see if you've missed a dose. Work with your doctor to decide on the medication plan that helps you the most. Clinical trials of Paxil produced several cautions: Treating bipolar disorder with Paxil (paroxetine) is considered an "off label" use of the drug, but doctors do prescribe it sometimes for this purpose. Bipolar disorder is a chronic mental health condition that causes people to experience periods of mania (racing and grandiose thoughts, increased physical and mental energy. Nov 8, - Learn about medications commonly used for bipolar disorder, how they work and possible side effects. Mar 10, - However, distressing side effects for drugs used in treatment of bipolar disorder is so common, and serious issues in the use of drug treatment, so common, that it is of serious Paxil is an SSRI of special note for those with bipolar disorder, or who might be depressed and have a tendency towards mania. Nov 8, - Once symptoms are under control, you'll receive maintenance treatment to reduce the risk of relapse. Maintenance treatment also reduces the chance of minor mood swings developing into mania or depression. Several types of medications are used to treat bipolar disorder. These include mood stabilizers. However, the recently published results from the STEP-BD project found no benefit to adding an antidepressant (paroxetine or bupropion; n =) compared with placebo (n =) to a mood stabilizer in a large naturalistic sample of patients with bipolar I and II disorders. This intriguing finding certainly questions whether. Serotonin reuptake inhibitors (SRIs), the drugs most commonly used for the treatment of OCD, can destabilize bipolar disorder by causing manic episodes or a the one that has been studied most extensively in bipolar depression (and that is associated with at least a relatively low rate of immediate mania) is paroxetine. Oct 20, - 1 Answer - Posted in: paxil, bipolar disorder, anxiety - Answer: About the time you would expect the Paxil to start working for depression. However, do not expect medication alone to solve all your problems. There are plenty of other steps you can take to manage your symptoms and reduce the amount of medication required. Medication is most effective when used in combination with other bipolar disorder treatments, including therapy, self-help coping. Read about Paroxetine, a medication that works in the brain to treat major depressive disorder. People with bipolar disorder who take antidepressants may be at risk for "switching" from depression into mania. . Paroxetine should not be taken with or within two weeks of taking monoamine oxidase inhibitors (MAOIs). Mar 14, - Paxil (generic paroxetine) is an antidepressant medication that may be prescribed for bipolar and major depression as well as several anxiety disorders.